

Abstract Submitted
for the Thirteenth International Conference on Laser
Interactions and Related Plasma Phenomena (LIRPP)
April 13-18, 1997

Absorption of Ultrashort Laser Pulses by Solid Targets
Heated Rapidly to Temperatures 1—1000 eV.* D. F. PRICE, R.
M. MORE, R. E. STEWART, G. GUETHLEIN, D. GOLD, R. S.
WALLING, W. E. WHITE, Lawrence Livermore National
Laboratory — We have measured the absorption of high-
contrast ultrashort pulses on a variety of solid targets over an
intensity range of 10^{13} to 10^{18} W/cm². Our measurements
give an experimental determination of the target energy
content and an indirect measure of dense plasma electrical
properties. The data include total absorption as a function of
intensity and angle of incidence, and absorption on metal
targets with oxide overlayers of varying thickness. Straightforward theoretical calculations accurately reproduce
the behavior of aluminum targets, while the other materials
show signs of an additional absorption mechanism. We have
characterized this additional absorption for Al₂O₃ targets. At
high intensity all target materials reach a "universal plasma
mirror" state and reflect about 90% of the incident light.
*Supported in part under the auspices of the U.S. Department
of Energy by Lawrence Livermore National Laboratory under
Contract No. W-7405-ENG-48.

Submitted by

Richard M. More
Lawrence Livermore National Lab
P. O. Box 808, L-041
Livermore, CA 94550